

PROCEDURE SPECIFIC WRIST CPM THERAPY

OPEN REDUCTION

INTERNAL FIXATION^{3,4,5,6,8,12,13,14,15,20,21}

Set-Up

- CPM should be initiated within 24 hours postoperatively.^{8,11}
- If possible the patient should be instructed on wrist CPM use preoperatively to improve compliance.⁸
- The W2 wrist CPM is recommended for night use. The Maestra wrist CPM is recommended if an end range stretch is desired.

Wearing Schedule

- The CPM is used for 4-8 weeks, or longer, depending on the condition and specific physician PROM standard of practice.^{5,6,8}
- The CPM device initially is used 8-20 hours per day or as tolerated pain free (week one).⁸
- Usage per day may be decreased as swelling decreases. Daily use should increase if stiffness persists.⁸
- Duration of use is 4-6 weeks or as needed.⁸

PROM Goals

- Initially the CPM is set up through available ROM as determined by the physician.⁸
- “Full ROM attained during surgery may be less during the first 2-3 weeks due to edema.”⁸
- The patient increases or maintains ROM as tolerated, pain free to meet ROM goals.⁸

Note: Specific PROM and usage parameters determined by the physician.



W2 Wrist CPM



Maestra Wrist CPM

REFERENCES:

1. Chigey LK, Dell FC, Muniz RE. Continuous Passive Motion of the Wrist Following Arthroscopy and Ganglionectomy as means of influencing the Recovery of Optimal Wrist Motion. Study presented at ASHT Meeting Jan-March, 1990.
2. Daane, M: (2002) [Evaluation of CPM insurance coverage]. Unpublished raw data.
3. Hareau, J PT, OTR. What Makes Treatment for Reflex Sympathetic Dystrophy Successful? *J. Hand Ther.* 9: 367-370, Oct. - Dec., 1996.
4. Herbert TJ: *The fractured scaphoid*, St Louis, 1990, Quarterly Medical Publishing.
5. Herbert TJ, Fischer WE, Leicester AW: The Herbert bone screw: a ten year perspective, *J Hand Surg* 17B:415, 1992.
6. Herbert TJ, Fischer WE: Management of the fractured scaphoid using a new bone screw, *J Bone Joint Surg* 66:114, 1984.
7. Kaczander B: The podiatric application of continuous passive motion, a preliminary report. *J. Am. Pod. Med. Assoc.* 81: 12, 1991.
8. LaStayo PC, Cass R: Continuous passive motion for the upper extremity: why, when, and how. In Hunter JM, et al, editors: *Rehabilitation of the Hand*, ed 5 St Louis, 2002, Mosby.
9. Mooney V, Stills M: Continuous passive motion with joint fractures and infections, *Orthop Clin North Am* 18:1, 1987.
10. O'Driscoll SW, Giori NJ: Continuous passive motion (CPM): Theory and principles of clinical application, *J Rehab Res Dev* 37: 179, 2000.
11. Osterman AL, Bora FW, Skirven T: The use of continuous passive motion in hand rehabilitation. Proceedings of the 42nd annual meeting of the American Society for Surgery of the Hand. San Antonio, 1987.
12. Salter RB: The prevention of arthritis through the preservation of cartilage, Royal College Lecture, *J Can Assoc Radiol* 32:5, 1981.
13. Salter RB: The effect of motion on regeneration of cartilage. Proceedings of the International Workshop on Rehabilitation of Articular Joints by Biological Resurfacing, Dallas, November 15-17, 1979.
14. Salter RB: The healing of bone and cartilage in intraarticular fractures with continuous passive motion. Transactions of the twenty-fourth annual meeting of the Orthopaedic Research Society, vol 3, February 1978.
15. Salter RB, Bell RS: The effect of continuous passive motion in the healing of partial thickness lacerations of the patellar tendon of the rabbit, *Ann Royal Coll Phys Surg Can* 14:209, 1981.
16. Salter RB, O'Driscoll SW: The effects of continuous passive motion on repair of full thickness defects in a joint surface with autogenous osteoperiosteal grafts. *Ann Royal Coll Phys Surg Can* 16:360, 1983 (abstract).
17. Salter RB, Minster RR: The effects of continuous passive motion on a semitendinous tenodesis in the rabbit knee, *Proc Orthop Res Soc* 7:225, 1982.
18. Salter RB, Bell RS, Keely F: The protective effect of continuous passive motion on living articular cartilage in acute septic arthritis, an experimental investigation in the rabbit, *Clin Orthop Rel Res* 159:223, 1981.
19. Salter RB, Harris DJ, Bogoch F: Further studies in continuous passive motion. *Orthop Trans* 212, 1978 (abstract).
20. Salter RB, Wong DA, Keeley FW: Collagen typing of early repair tissue in healing articular cartilage: an experimental study in the rabbit. *Orthop Trans* 4:397, 1980.
21. Schatzker J, Tile M: *The rational of operative fracture care*, New York, 1987, Springer-Verlag.